

Diagonalization of compact operators in Hilbert modules over C^* -algebras of real rank zero

V. M. Manuilov

22 January 1995

Abstract

It is known that the classical Hilbert–Schmidt theorem can be generalized to the case of compact operators in Hilbert \mathcal{A} -modules $\mathcal{H}_{\mathcal{A}}^*$ over a W^* -algebra of finite type, i.e. compact operators in $\mathcal{H}_{\mathcal{A}}^*$ under slight restrictions can be diagonalized over \mathcal{A} . We show that if \mathcal{B} is a weakly dense C^* -subalgebra of real rank zero in \mathcal{A} with some additional property then the natural extension of a compact operator from $\mathcal{H}_{\mathcal{B}}$ to $\mathcal{H}_{\mathcal{A}}^* \supset \mathcal{H}_{\mathcal{B}}$ can be diagonalized with diagonal entries being from the C^* -algebra \mathcal{B} .

1 Introduction

Let \mathcal{A} be a C^* -algebra. We consider Hilbert \mathcal{A} -modules over \mathcal{A} [13], i.e. (right) \mathcal{A} -modules \mathcal{M} together with an \mathcal{A} -valued inner product $\langle \cdot, \cdot \rangle : \mathcal{M} \times \mathcal{M} \rightarrow \mathcal{A}$ satisfying the following conditions:

- i) $\langle x, x \rangle \geq 0$ for every $x \in \mathcal{M}$ and $\langle x, x \rangle = 0$ iff $x = 0$,
- ii) $\langle x, y \rangle = \langle y, x \rangle^*$ for every $x, y \in \mathcal{M}$,
- iii) $\langle \cdot, \cdot \rangle$ is \mathcal{A} -linear in the second argument,
- iv) \mathcal{M} is complete with respect to the norm $\|x\|^2 = \|\langle x, x \rangle\|_{\mathcal{A}}$.

By $\mathcal{M}^* = \text{Hom}_{\mathcal{A}}(\mathcal{M}; \mathcal{A})$ we denote the \mathcal{A} -module dual to \mathcal{M} . Let $\mathcal{H}_{\mathcal{A}}$ be a right Hilbert \mathcal{A} -module of sequences $a = (a_k)$, $a_k \in \mathcal{A}$, $k \in \mathbf{N}$ such that the series $\sum a_k^* a_k$ converges in \mathcal{A} in norm with the standard basis $\{e_k\}$ and let $L_n(\mathcal{A}) \subset \mathcal{H}_{\mathcal{A}}$ be a submodule generated by the elements e_1, \dots, e_n of the basis. An inner \mathcal{A} -valued product on module $\mathcal{H}_{\mathcal{A}}$ is given by $\langle x, y \rangle = \sum x_k^* y_k$ for $x, y \in \mathcal{H}_{\mathcal{A}}$. A bounded operator $\mathcal{K} : \mathcal{H}_{\mathcal{A}} \rightarrow \mathcal{H}_{\mathcal{A}}$ is called compact [7] [8], if it possesses an adjoint operator and lies in the norm closure of the linear span of operators of the form $\theta_{x,y}$, $\theta_{x,y}(z) = x\langle y, z \rangle$, $x, y, z \in \mathcal{H}_{\mathcal{A}}$. From now on we

suppose that the compact operator \mathcal{K} is strictly positive, i.e. operator $\langle \mathcal{K}x, x \rangle$ is positive in \mathcal{A} and $\text{Ker } \mathcal{K} = 0$. It is known [14] that in the case when \mathcal{A} is a W^* -algebra the inner product can be naturally prolonged to the dual module $\mathcal{H}_{\mathcal{A}}^*$.

Definition 1.1. Let \mathcal{A} be a W^* -algebra. We call an operator \mathcal{K} *diagonalizable* if there exist a set $\{x_i\}$ of elements in $\mathcal{H}_{\mathcal{A}}^*$ and a set of operators $\lambda \in \mathcal{A}$ such that

- i) $\{x_i\}$ is orthonormal, $\langle x_i, x_j \rangle = \delta_{ij}$,
- ii) $\mathcal{H}_{\mathcal{A}}^*$ coincides with the \mathcal{A} -module \mathcal{M}^* dual to the module \mathcal{M} generated by the set $\{x_i\}$,
- iii) $\mathcal{K}x_i = x_i\lambda_i$,
- iv) for any unitaries $u_i, u_{i+1} \in \mathcal{A}$ we have an operator inequality

$$u_i^* \lambda_i u_i \geq u_{i+1}^* \lambda_{i+1} u_{i+1}. \quad (1.1)$$

We call the elements x_i “*eigenvectors*” and the operators λ_i “*eigenvalues*” for the operator \mathcal{K} . It must be noticed that the “eigenvectors” and “eigenvalues” are defined not uniquely.

The problem of diagonalizing operators in Hilbert modules was initiated by R. V. Kadison in [6] and was studied in different settings in [5],[11],[4],[15] etc. In [9],[10] we have proved the following

Theorem 1.2. *If \mathcal{A} is a finite σ -finite W^* -algebra then a compact strictly positive operator \mathcal{K} can be diagonalized and its “eigenvalues” are defined uniquely up to unitary equivalence.*

It is well known that in the commutative case, i.e. for $\mathcal{C} = C(X)$ being a commutative C^* -algebra, compact operators cannot be diagonalized inside $\mathcal{H}_{\mathcal{C}}$ but it becomes possible if we pass to a bigger module over a bigger W^* -algebra $L^\infty(X) \supset \mathcal{C}$. It leads us to the following

Definition 1.3. Let \mathcal{C} be a C^* -algebra admitting a weakly dense inclusion in a finite σ -finite W^* -algebra \mathcal{A} and let \mathcal{K} be a compact strictly positive operator in $\mathcal{H}_{\mathcal{C}}$. We can naturally extend \mathcal{K} to the bigger module $H_{\mathcal{A}}^*$ where it will remain compact and strictly positive and by the theorem 1.2 it can be diagonalized in this module. We call a C^* -algebra \mathcal{C} admitting *weak diagonalization* if the diagonal entries for any \mathcal{K} in $\mathcal{H}_{\mathcal{A}}^*$ can be taken from \mathcal{C} instead of \mathcal{A} .

Problem. Describe the class of C^* -algebras admitting weak diagonalization.

Throughout this paper we denote by \mathcal{A} a finite σ -finite W^* -algebra. Denote by $\mathcal{Z} = C(\mathcal{Z})$ the center of \mathcal{A} and by T the standard exact center-valued trace

defined on \mathcal{A} , $T(\mathbf{1}) = 1$. Suppose that for a C^* -subalgebra \mathcal{B} of \mathcal{A} the following condition holds:

- (*) for any two projections $p, q \in \mathcal{B}$ there exist in \mathcal{B} equivalent (in \mathcal{B}) projections $r_p \sim r_q$, $r_p \leq p$, $r_q \leq q$ such that $T(r_p) = T(r_q) = \min\{T(p)(z), T(q)(z)\}$, $z \in Z$.

The purpose of this paper is to show that the class of C^* -algebras admitting weak diagonalization contains real rank zero weakly dense C^* -subalgebras of finite σ -finite W^* -algebras with the property (*). Recall that real rank zero ($RR(\mathcal{B}) = 0$) means [2] that every selfadjoint operator in \mathcal{B} can be approximated by operators with finite spectrum, i.e. having the form $\sum \alpha_i p_i$, where $p_i \in \mathcal{B}$ are selfadjoint mutually orthogonal projections and $\alpha_i \in \mathbf{R}$. By [2] we have in this case also $RR(\text{End}_{\mathcal{B}}(L_n(\mathcal{B}))) = 0$.

2 Continuity of “eigenvalues”

For the further we need to establish some continuity properties of the “eigenvalues” of compact operators in modules over W^* -algebras.

Lemma 2.1. *Let $\mathcal{K}_1 = \sum \alpha_l^{(1)} P_l^{(1)}$, $\mathcal{K}_2 = \sum \alpha_l^{(2)} P_l^{(2)}$ be strictly positive operators in $L_n(\mathcal{A})$ with finite spectrum and let $\|\mathcal{K}_1 - \mathcal{K}_2\| < \varepsilon$. Then*

- i) *one can find a unitary U in $L_n(\mathcal{A})$ such that it maps the “eigenvectors” of \mathcal{K}_2 to the “eigenvectors” of \mathcal{K}_1 and $\|U^* \mathcal{K}_1 U - \mathcal{K}_2\| < \varepsilon$,*
- ii) *“eigenvalues” $\{\lambda_i^{(r)}\}$ of operators \mathcal{K}_r ($r = 1, 2$) can be chosen in such a way that $\|\lambda_i^{(1)} - \lambda_i^{(2)}\| < \varepsilon$.*

Proof. As the algebra \mathcal{A} can be decomposed into a direct integral of finite factors, so it is sufficient to prove the lemma for the case when \mathcal{A} is a type II_1 factor (for type I_n factors lemma is trivial). Denote by $E_{\mathcal{K}}(\lambda)$ the spectral projection for the operator \mathcal{K} corresponding to the set $(-\infty, \lambda)$. If τ is an exact finite trace on \mathcal{A} , it can be prolonged to the (infinite) trace $\bar{\tau} = \text{tr} \otimes \tau$ on the algebra $\text{End}_{\mathcal{A}}(\mathcal{H}_{\mathcal{A}})$ and to the finite trace on a lesser algebra $\text{End}_{\mathcal{A}}(L_n(\mathcal{A}))$ where we have $\bar{\tau}(\mathbf{1}) = n$. Put

$$\varepsilon_{\mathcal{K}}(\alpha) = \inf_{\bar{\tau}(E_{\mathcal{K}}(\lambda)) \geq \alpha} \lambda, \quad 0 \leq \alpha \leq n.$$

As it is shown in [12] (the continuous minimax principle) one has

$$\varepsilon_{\mathcal{K}}(\alpha) = \inf_{P \in \mathcal{P}, \bar{\tau}(P) \geq \alpha} \left\{ \sup_{\xi \in \text{Im } P, \|\xi\|=1} (\mathcal{K}\xi, \xi) \right\}, \quad (2.1)$$

where (\cdot, \cdot) denotes an inner product in a Hilbert space where the algebra $\text{End}_{\mathcal{A}}(L_n(\mathcal{A}))$ is represented and \mathcal{P} denotes the set of projections in $\text{End}_{\mathcal{A}}(L_n(\mathcal{A}))$. It follows from (2.1) that if $\|\mathcal{K}_1 - \mathcal{K}_2\| < \varepsilon$, then

$$|\varepsilon_{\mathcal{K}_1}(\alpha) - \varepsilon_{\mathcal{K}_2}(\alpha)| < \varepsilon. \quad (2.2)$$

Let $Q_i^{(r)}$ be projections on the “eigenvectors” $x_i^{(r)}$ of the operators \mathcal{K}_r , corresponding to the maximal “eigenvalues” $\lambda_i^{(r)}$, $\bar{\tau}(Q_i^{(r)}) = 1$. For two divisions $\{P_l^{(1)}, Q_i^{(1)}\}$ and $\{P_l^{(2)}, Q_i^{(2)}\}$ of unity given by decompositions of \mathcal{K}_1 and \mathcal{K}_2 we can construct a finer division of unity. By [16] there exist sets of mutually orthogonal projections $R_m^{(r)} \in \text{End}_{\mathcal{A}}(L_n(\mathcal{A}))$ such that

$$i) \quad \bigoplus_m R_m^{(r)} = 1,$$

$$ii) \quad \bar{\tau}(R_m^{(1)}) = \bar{\tau}(R_m^{(2)}),$$

$$iii) \quad \text{for every } m \text{ we have } R_m^{(r)} \leq Q_i^{(r)} \text{ or } R_m^{(r)} \leq P_j^{(r)} \text{ for some } i \text{ or } j.$$

Then (after renumbering) one can write the operators \mathcal{K}_r in the form $\mathcal{K}_r = \sum \alpha_m^{(r)} R_m^{(r)}$ with $\alpha_1^{(r)} \leq \alpha_2^{(r)} \leq \dots$, $\alpha_m^{(r)} \in \mathbf{R}$. It makes possible to define a unitary $U : L_n(\mathcal{A}) \rightarrow L_n(\mathcal{A})$ such that

$$U(\text{Im } R_m^{(2)}) = \text{Im } R_m^{(1)}, \quad (2.3)$$

hence $U(\text{Im } Q_i^{(2)}) = \text{Im } Q_i^{(1)}$ so U maps the \mathcal{A} -modules generated by the “eigenvectors” $x_i^{(2)}$ into the modules generated by $x_i^{(1)}$, hence $Ux_i^{(2)} = x_i^{(1)} \cdot u_i = \bar{x}_i^{(1)}$ for some unitaries $u_i \in \mathcal{A}$. Put

$$n(\alpha) = \min\{n | \bar{\tau}(\bigoplus_{m \geq n} R_m^{(r)}) \geq \alpha\}.$$

Then $\varepsilon_{\mathcal{K}_r(\alpha)} = \alpha_{n(\alpha)}^{(r)}$ and it follows from (2.2) that $|\alpha_{n(\alpha)}^{(1)} - \alpha_{n(\alpha)}^{(2)}| < \varepsilon$. But changing α we obtain that

$$|\alpha_m^{(1)} - \alpha_m^{(2)}| < \varepsilon \quad (2.4)$$

for all m . Taking $\alpha = 1$ (then $i = 1$) we have

$$\mathcal{K}_r|_{\text{Im } Q_1^{(r)}} = \Lambda_1^{(r)} = \sum_{m \geq n(1)} \alpha_m^{(r)} P_m^{(r)}.$$

From (2.3) and (2.4) we conclude that

$$\|U^* \Lambda_1^{(1)} U - \Lambda_1^{(2)}\| = \left\| \sum_{m \geq n(1)} (\alpha_m^{(1)} - \alpha_m^{(2)}) P_m^{(2)} \right\| \leq \varepsilon \left\| \bigoplus_{m \geq n(1)} P_m^{(2)} \right\| = \varepsilon. \quad (2.5)$$

Choosing appropriate $\lambda_1^{(r)}$ to satisfy the conditions $\Lambda_1^{(1)}\bar{x}_1^{(1)} = \bar{x}_1^{(1)}\lambda_1^{(1)}$ and $\Lambda_1^{(2)}x_1^{(2)} = x_1^{(2)}\lambda_1^{(2)}$ we obtain the estimate

$$\|\lambda_1^{(1)} - \lambda_1^{(2)}\| < \varepsilon. \quad (2.6)$$

By the same way estimates (2.5), (2.6) can be obtained for all i and it proves the lemma. •

Corollary 2.2. *Let $\mathcal{K}_r : \mathcal{H}_{\mathcal{A}} \rightarrow \mathcal{H}_{\mathcal{A}}$, $r = 1, 2$ be compact strictly positive operators and let $\|\mathcal{K}_1 - \mathcal{K}_2\| < \varepsilon$. Then*

- i) one can find a unitary U in $\mathcal{H}_{\mathcal{A}}^*$ such that it maps the “eigenvectors” of \mathcal{K}_2 to the “eigenvectors” of \mathcal{K}_1 and $\|U^*\mathcal{K}_1U - \mathcal{K}_2\| < \varepsilon$,*
- ii) “eigenvalues” $\{\lambda_i^{(r)}\}$ of operators \mathcal{K}_r ($r = 1, 2$) can be chosen in such a way that $\|\lambda_i^{(1)} - \lambda_i^{(2)}\| < \varepsilon$.*

Proof. Let $L_n^{(r)}(\mathcal{A}) \in \mathcal{H}_{\mathcal{A}}^*$ denotes the Hilbert submodule generated by the first n “eigenvectors” of the operator \mathcal{K}_r , $L_n^{(r)}(\mathcal{A}) \cong L_n(\mathcal{A})$. It was shown in [10] that the orthogonal complement to such submodule is isomorphic to $\mathcal{H}_{\mathcal{A}}^*$ and the norm of restriction of compact operator \mathcal{K}_r on the orthogonal complement to $L_n^{(r)}(\mathcal{A})$ in $\mathcal{H}_{\mathcal{A}}^*$ tends to zero, henceforth it is sufficient to consider only the case of operators in $L_n(\mathcal{A})$ and there one can approximate these operators by operators with finite spectrum. •

3 Case of $RR(\mathcal{B}) = 0$

In this section we show that C^* -algebras of real rank zero with the property (*) admit weak diagonalization.

Theorem 3.1. *Let \mathcal{B} be a weakly dense C^* -subalgebra in \mathcal{A} with the property (*) and let $RR(\mathcal{B}) = 0$. If \mathcal{K} is a compact strictly positive operator in the \mathcal{B} -module $\mathcal{H}_{\mathcal{B}}$ then the “eigenvalues” $\{\lambda_i\}$ of diagonalization of the natural prolongation of \mathcal{K} to the \mathcal{A} -module $\mathcal{H}_{\mathcal{A}}^*$ can be chosen in a way that $\lambda_i \in \mathcal{B}$ would hold.*

Proof is based on the results of S. Zhang [17]. By [2],[17] the operator \mathcal{K} can be approximated by operators $\mathcal{K}_n \in \text{End}_{\mathcal{B}}(L_n(\mathcal{B}))$ with finite spectrum. By [17], corollary 3.5 there exist such unitaries $U_n \in \text{End}_{\mathcal{B}}(L_n(\mathcal{B}))$ that the operators

$$U_n^*\mathcal{K}_nU_n = \begin{pmatrix} \lambda_1^{(n)} & & 0 \\ & \ddots & \\ 0 & & \lambda_n^{(n)} \end{pmatrix}$$

are diagonal and $\lambda_i^{(n)} \in \mathcal{B}$ are operators with finite spectrum. Show that due to the property (*) by an appropriate choice of such U_n one can make the condition (1.1) valid for “eigenvalues” $\{\lambda_i^{(n)}\}$. Let $\lambda_a = \sum \alpha_k q_k$, $\lambda_b = \sum \beta_l r_l$ where $q_k, r_l \in \mathcal{B}$ are projections and suppose that $a < b$ but for some m and n inequality $\beta_m > \alpha_n$ holds. Using the possibility to diagonalize projections [17] we can find projections $s_l \in \mathcal{B}$ equivalent to r_l and such that $s_l = \oplus_k s_k^{(l)}$ and $s_k^{(l)} \leq q_k$. Then put

$$\lambda'_a = \sum_{k \neq n} \alpha_k q_k \oplus \sum_{l \neq m} \alpha_n s_n^{(l)} \oplus \beta_n s_n^{(m)},$$

$$\lambda'_b = \sum_{l \neq m} \beta_l s_l \oplus \sum_{k \neq n} \beta_k s_k^{(m)} \oplus \alpha_n s_n^{(m)}$$

and notice that the operators $\begin{pmatrix} \lambda_1 & & 0 \\ & \ddots & \\ 0 & & \lambda_n \end{pmatrix}$ and $\begin{pmatrix} \lambda'_1 & & 0 \\ & \ddots & \\ 0 & & \lambda'_n \end{pmatrix}$ are unitarily equivalent. After repeating this procedure for all cases when $\beta_l > \alpha_k$ we obtain validity of (1.1) for λ'_a and λ'_b . By the same way we can order all “eigenvalues” of \mathcal{K}_n remaining in \mathcal{B} . But by the property (*) if $\|\mathcal{K}_n - \mathcal{K}_{n-1}\| < \varepsilon_n$ then one can find such unitaries $u_{i,n}$ in \mathcal{B} that

$$\|u_{i,n}^* \lambda_i^{(n)} u_{i,n} - \lambda_i^{(n-1)}\| < \varepsilon_n \quad (3.1)$$

. Then $u_{i,n}^* \lambda_i^{(n)} u_{i,n} \in \mathcal{B}$. Taking a subsequence of $\{\mathcal{K}_n\}$ if necessary we can take in (3.1) $\varepsilon_n = \frac{1}{2^n}$. Then the sequence

$$\bar{\lambda}_i^{(1)} = \lambda_i^{(1)}, \bar{\lambda}_i^{(2)} = u_{i,2}^* \lambda_i^{(2)} u_{i,2}, \bar{\lambda}_i^{(3)} = u_{i,3}^* u_{i,2}^* \lambda_i^{(3)} u_{i,2} u_{i,3}, \dots$$

is fundamental in \mathcal{B} . Denote its limit by $\bar{\lambda}_i \in \mathcal{B}$. By the corollary 2.2 for all \mathcal{K}_n we can find unitaries U_n which map the first n “eigenvectors” of \mathcal{K} to “eigenvectors” of \mathcal{K}_n . Put $\mathcal{K}'_n = U_n^* \mathcal{K}_n U_n \in \text{End}_{\mathcal{A}}(\mathcal{H}_{\mathcal{A}}^*)$. Then we have

$$\mathcal{K}'_n x_i = x_i \bar{\lambda}_i^{(n)} \quad (3.2)$$

and $\|\mathcal{K}'_n - \mathcal{K}\| \rightarrow 0$. Taking limit in (3.2) we obtain $\mathcal{K} x_i = x_i \bar{\lambda}_i$, hence $\bar{\lambda}_i$ are “eigenvalues” of \mathcal{K} . •

Notice that the condition (*) is necessary for a C^* -algebra to have the weak diagonalization property. Indeed if \mathcal{K} is a direct sum of two projections, $\mathcal{K} = \begin{pmatrix} p & 0 \\ 0 & q \end{pmatrix}$ then the “eigenvalues” of \mathcal{K} can be ordered only if the “common part” of $1 - p$ and q lies in \mathcal{B} .

Remark. In the case of C^* -algebras A_θ of irrational rotation one has $RR(A_\theta) = 0$ (cf [3]) and the property (*) is valid, so the theorem 3.1 gives

the answer to the problem of [10] where we have considered the Schrödinger operator in magnetic field with irrational magnetic flow. It is known that this operator can be viewed as an operator acting in a Hilbert A_θ -module. As we can imbed A_θ in a type II_1 factor \mathcal{A} as a weakly dense subalgebra [1] so we can diagonalize this operator in a Hilbert \mathcal{A} -module. The present paper shows that the “eigenvalues” of this operator can be chosen to be elements of A_θ . So this situation is a noncommutative analogue of the case $\theta = 1$ when the corresponding operator can be diagonalized over W^* -algebra $L^\infty(\mathbf{T}^2)$ but the diagonal elements lie in a lesser C^* -algebra $C(\mathbf{T}^2)$. Notice that in case of rational θ this operator is also diagonalizable.

Acknowledgement. This work was partially supported by the Russian Foundation for Fundamental Research (grant N 94-01-00108-a) and the International Science Foundation (grant N MGM000). I am indebted to M. Frank, A. A. Irmatov, A. S. Mishchenko and E. V. Troitsky for helpful discussions.

References

- [1] BRENKEN B.: Representations and automorphisms of the irrational rotation algebra. *Pacif. J. Math.* **111** (1984), 257 – 282.
- [2] BROWN L.G., PEDERSEN G.K.: C^* -algebras of real rank zero. *J. Funct. Anal.* **99** (1991), 131 – 149.
- [3] CHOI M.-D., ELLIOT G.A.: Density of self-adjoint elements with finite spectrum in an irrational rotation C^* -algebra. *Math. Scand.* **67** (1990), 73 – 86.
- [4] FRANK M., MANUILOV V.M.: Diagonalizing “compact” operators on Hilbert W^* -modules. *Zeitschr. Anal. Anwendungen.* **14** (1995), 33 – 41.
- [5] GROVE K., PEDERSEN G.K.: Diagonalizing matrices over $C(X)$. *J. Funct. Anal.* **59** (1984), 64 – 89.
- [6] KADISON R.V.: Diagonalizing matrices. *Amer. J. Math.* **106** (1984), 1451 – 1468.
- [7] KASPAROV G.G.: Topological invariants of elliptic operators. I. K -homology. *Izv. Akad. Nauk SSSR. Ser. Mat.* **39** (1975), 796 – 838 (in Russian).
- [8] MISHCHENKO A.S., FOMENKO A.T.: The index of elliptic operators over C^* -algebras. *Izv. Akad. Nauk SSSR. Ser. Mat.* **43** (1979), 831 – 859 (in Russian).
- [9] MANUILOV V.M.: Diagonalization of compact operators in Hilbert modules over W^* -algebras of finite type. *Usp. Mat. Nauk* **49** (1994), No 2, 159 - 160 (in Russian).
- [10] MANUILOV V.M.: Diagonalization of compact operators in Hilbert modules over finite W^* -algebras. to appear in *Annals of Global Anal. Geom.* **13** (1995).

- [11] MURPHY Q.J.: Diagonality in C^* -algebras. *Math. Zeitschr.* **199** (1990), 279 – 284.
- [12] MURRAY F.J., VON NEUMANN J.: On rings of operators. *Ann. Math.* **37** (1936), 116 – 229.
- [13] PASCHKE W.L.: Inner product modules over B^* -algebras. *Trans. Amer. Math. Soc.* **182** (1973), 443 – 468.
- [14] PASCHKE W.L.: The double B -dual of an inner product module over a C^* -algebra. *Canad. J. Math.* **26** (1974), 1272 – 1280.
- [15] SUNDER V.S., THOMSEN K.: Unitary orbits of selfadjoints in some C^* -algebras. *Houston J. Math.* **18** (1992), 127 – 137.
- [16] TAKESAKI M.: Theory of operator algebras,1. New-York – Heidelberg – Berlin: Springer Verlag, 1979.
- [17] ZHANG S.: Diagonalizing projections. *Pacif. J. Math.* **145**(1990), 181 – 200.

V. M. Manuilov
 Moscow State University
 Moscow, 119899, Russia
 E-mail:manuilov@mech.math.msu.su